



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Doassansia ranunculina, n. sp.—Spots light brown, 2–4^{mm} in diameter, papillate. Sori aggregated or scattered, usually spherical, 100–200 μ in diameter, developing in either the parenchyma or the palisade layer of the leaf or in any position in the petiole between the epidermis and the fibro-vascular bundles. Spores crowded, filling the sorus, spherical or polygonal, 6–10 μ in diameter. Cortex of one layer of cells which are more or less quadrangular in section, 10–15 \times 8–12 μ . Spores germinating in position. Promycelium 3–4 μ in diameter, vacuolate. Primary sporidia 6–8 in a whorl on the end of the promycelium, fusiform, vacuolate, 12–20 \times 2–2.5 μ . Conjugation by means of a large apical tube both in position and after becoming free. A filament 60–90 μ long is then formed from which the secondary sporidia are abstricted in basipetal succession. Secondary sporidia 12–15 \times 2.5–3 μ . In material which had germinated and produced the filaments in the field the secondary sporidia conjugated by means of connecting tubes. In many of the specimens collected germination had occurred and the filaments protruded through the cleft cortex and epidermis.

In the leaves and occasionally the petioles of *Ranunculus multifidus* Pursh, in swampy places which had become dry during the summer. Racine, Wisconsin.

Material has been prepared for the distribution of these species in Ellis and Everhart's North American Fungi.—J. J. DAVIS, *Racine, Wisconsin*.

Ruled slides.—For several years I have made much use of ruled slides with a stage microscope. It is so easy thus to measure any thing you dissect, that I wonder they are not in more general use. Those I use are ruled in squares, 10^{mm} each way, in the middle of a slide which ought to be three inches long, and wider than the ordinary slide to prevent getting mixed up with them. The ruling ought to be deep enough so that water will not temporarily make it impossible or very difficult to see the lines. Good ruled slides for the use above named ought to be in the market at moderate prices.—W. J. BEAL, *Agricultural College, Mich.*